

CERN-PH/2006-118
 PNPI/1000/06
 LAPTH/1155/06

To our Friend Tatiana Fabergé

We can only find new ideas about
 our World and suggest ways of
 checking them in experiments.
 But whether they are right or
 wrong - only GOD knows.
 Bruno Pontecorvo, Dubna, 1981

Geometry of the Majorana Neutrino and New Symmetries

G. G. Volkov ¹

Abstract

Experimental observation of Majorana fermion matter gives a new impetus to the understanding of the Lorentz symmetry and its extension, the geometrical properties of the ambient space-time structure, matter-antimatter symmetry and some new ways to understand the baryo-genesis problem in cosmology. Based on the primordial Majorana fermion matter assumption, we discuss a possibility to solve the baryo-genesis problem through the the Majorana-Diraco genesis in which we have a chance to understand creation of Q_{em} charge and its conservation in our $D = 1 + 3$ Universe after the Big Bang . In the Majorana-Diraco genesis approach there appears a possibility to check the idea of the proton and electron non-stability on the very low energy scale. In particle physics and in our space-time geometry, the Majorana nature of the neutrino can be related to new types of symmetries which are lying beyond the binary Cartan-Killing-Lie algebras/superalgebras. This can just support a conjecture about the non-completeness of the SM in terms of binary Cartan-Killing-Lie symmetries/supersymmetries. As one of the very important applications of such new ternary symmetries could be related with explanation of the nature of the three families and three colour symmetry. Based on the Majorana-Diraco genesis the Majorana neutrino can directly indicate the existence of a new extra-dimensional geometry (new cycle) and thanks to new ternary space-time symmetries, could lead at high energies to the unextraordinary phenomenological consequences.

¹On leave from PNPI, Russia

We would like to discuss how the new space-time properties of the neutrino, namely its Majorana properties, may be an indication of extra uncompactified space-time dimensions. The existence of the Majorana neutrino gives new ideas about the possible composite structure and form of electromagnetism in our Universe through the relation between Dirac and Majorana fermions.

In this line of thought the experimental observation of the Majorana nature of the neutrino by the HEIDELBERG–MOSCOW experiment $0\nu\beta\beta$ decay [1, 2, 3] may have very important significance for our attempts to build a theory that encompasses the Standard Model.

Firstly, one can say that the discovery of $0\nu\beta\beta$ decay is the indication for Majorana fermionic matter in our Universe as was expected not just in some extension of the Standard Model, such as $L - -R$ models or Grand Unified Theory, but also in all supersymmetry, supergravity, superstring approaches.

Secondly, the Majorana fermions can have a much closer link with the geometrical properties of our space-time and, and most important, the Majorana neutrino can give the first evidence for new uncompactified extra-dimensions space with $D \geq 1 + 3$ and, correspondingly, the chance to observe the new space-time and internal symmetries.

Extra-dimensional cosmology offers new possibilities to solve the baryo-genesis problem which, correspondingly, can give a new universal mechanism of the proton and electron non-stability. By universality we mean that proton and electron decays are related with the same dynamics. Also this new dynamics can change our representations of dark-matter and dark-energy problems.

However, extra uncompactified dimensions make quantum field theories much less comfortable, since the power counting is worse [10]. A possible way out is to suppose that the propagator is more convergent than $1/p^2$, such a behaviour can be obtained if we consider, instead of binary symmetry algebra, algebras with higher order relations (*that is, instead of binary operations such as addition or product of 2 elements, one starts with composition laws involving at least n elements of the considered algebra, n -ary algebras*). For instance ternary symmetry could be related with membrane dynamics.

These exotic symmetries can be related to algebras which are based on the generalization of the Lie binary commutation relation, $[A, B]_2 = AB - BA$, by the ternary commutation relations: $[A, B, C]_3 = ABC + BCA + CAB - BAC - ACB - CBA$; they have already been discussed in physics [16, 11, 18, 19, 20, 24], in algebra [17], in CY_3 geometry [21, 22], [23], respectively. In [9] the ternary symmetries have been found through the ternary norm-division algebras of dimensions 3^n , *i.e.* ternary complex numbers ($n = 1$), ternary "quaternions" ($n = 2$), ternary "octonions" ($n = 3$). The ternary numbers directly give us the generalizations of the Abelian, $su(3)$ and $g(2)$ binary algebras [9].

The new symmetries beyond the well-known binary Lie algebras/superalgebras could allow us to build the renormalizable theories for space-time geometry with dimension $D > 4$. It seems very plausible that using such ternary symmetries will offer a real possibility to overcome the problems with quantization of membranes and could be a further progress beyond string/superstring theories. Also, these new ternary algebras could be related with some new supersymmetry approach [20]. Getting the renormalizable

quantum field theories in $D \geq 4$ space-time we access the point-like limits of the string and membrane theories for some extra dimension $6 \geq D \geq 4$.

The new, so low, mass-energy scale $\langle m \rangle_{ee} = 0.05 - 0.84 \text{ eV}$ (the best value being $0, 39 \text{ eV}$), $T_{1/2}^{0\nu} = (0.7 - 18.3) \times 10^{25} \text{ years}$ can indicate the discovery of a new space-time and of internal symmetries. We suppose that the ternary symmetry can be responsible for such low Majorana neutrino masses. Combining this with the neutrino oscillation experiments discovers an internal symmetry, leading to the mass relation

$$m_{\nu_e} = m_{\nu_\mu} = m_{\nu_\tau}, \text{ TERNARY RELATION} \quad (1)$$

which could be the first real indication of the existence of a ternary symmetry whose nature is lying beyond the ordinary Lie algebra. This new ternary symmetry could shed light on the "dark symmetry" of the SM:

$$N_{colour} = N_{family} = N_{dim} = 3. \quad (2)$$

This ternary mass relation reminds the binary Pauli-Luders relation between masses of particles and antiparticle following in $D = 3 + 1$ from CPT invariance and $D = 3 + 1$ Lorentz symmetry. We suppose that it could be followed a ternary $D = 6$ symmetry by generalized Lorentz symmetry and an analog of D=6 complete discrete symmetry in D=6. Thus we can conjecture such a relation just for the particles living in D=6, like Majorana neutrinos, ν_e, ν_μ, ν_τ . In this case the mechanism of oscillation for neutrinos can recall the mechanism of oscillation $K^0 - \bar{K}^0, B_d^0 - \bar{B}_d^0, \dots$. For this we should suggest a new interaction, which would mix the three neutrino states.

The first consequence is related an idea of complexity of all Dirac spinor-fields in the Standard Model apart from Majorana neutrino fields. The prediction of antimatter by Dirac in his relativistic equation and the further experimental confirmation of his idea were among the exciting discoveries in particle physics. But this discovery made a very intriguing puzzle for cosmology, which was not solved till now, 70-80 years after the discovery of antimatter.

The discovery of the Majorana nature neutrino among the set of all other Standard Model Dirac charged fermions prompts another dynamics of baryo-genesis, based on the composite dynamics of appearance of the fermion Dirac matter from simple real fermions like Majorana neutrino living in an extra-dimensional world?! The dynamics based on the extra-dimensional geometry could be beyond Sakharov's three conjectures. The fundamental conception of such an idea is related attempts to find a common mechanism of the creation of Q_{em} charge symmetry with baryo-genesis. We propose that such a mechanism must use a duality between the Q_{em} conservation and CPT invariance:

$$\text{CPT invariance} \quad \leftrightarrow \quad (Q_{em}) \text{ charge conservation}, \quad (3)$$

i.e. the invariance of CPT in $D = 4$ space-time means the conservation of the Q_{em} charge there, and opposite.

We could propose that primordially the Universe was neutral $Q_{em} = 0$, more over, there were absolutely no charged particles, in particularly no Dirac complex fermions. The Dirac complex fermions appeared in $D = 1 + 3$ "brane world" from real high-dimensional

fermions as a result of a dynamics having geometrical origin. In this case the part of a space-time where such process took place must have the total charge equal zero. We can propose that such higher dimensional space-time world has a symmetry which is more fundamental than the symmetry used in the Standard Model and Standard Cosmology Model, based on the Lie algebras and Lie superalgebras. We already have some candidates for such symmetries. They can be related with the n -ary generalizations ($n = 3, 4, \dots$) of Lie algebras and superalgebras. The geometry of such space-time and particle-states in this world can be determined by these new universal symmetries, for example, based on the unification of binary and ternary algebras. This could be a new way to construct a realistic Grand Unified Theory. Our proposal is that the Majorana neutrino could link with such neutral real fermions and therefore give us useful information about this high-dimensional world with $D > 4$. For terminology we can also use the word “brane world” which means that the Standard Model matter is “living” in the three-dimensional submanifold embedded into the higher-dimensional space (a confinement)[14].

Usually, it was suggested that only gravitons can penetrate into the bulk and only the neutrinos could interact with some bulk matter. Based on the discovery of Majorana fermions we accept the more general scenario in which the Majorana neutrinos can also be trapped in the bulk and some exotic real Majorana-like fermions from the bulk could interact weakly with Dirac fermions on the brane.

We can relate these real bulk fermions and neutrino to the possibility of creation of Dirac fermions and appearance of the Q_{em} charge symmetry in such world from some bulk real fermions. Such dynamics could lead to the simultaneous production of baryon and leptons after Big Bang by the unique mechanism leading to $Q_p + Q_e = 0$. In such a scenario the Majorana neutrinos properties should allow it to have a “memory” about this production-process?! We can study the physical models with extra dimensions through the new symmetries, which could help us to observe a duality between external and internal dynamical symmetries, *i.e.* to overcome the Coleman–Mandula no-go theorem. One of the ways in this direction is related to the search for generalizations of the principle of the theory of special relativity, generalizations of the Klein–Gordon or Dirac–Majorana equations [24]. There were some attempts to generalize the Dirac equation in the frames of some ternary Clifford algebra (see for example, [20]). The main question here is to get a “relativistic” invariant equation. The ternary norm-division algebras [9] gives the way to construct a generalization of the Lorentz symmetry [9] and, consequently, to get the invariant wave equations for the particles living in $D > 4$ space-time. Of course, the wave equations for the new “gauge” fields will contain the derivatives of degree more than 2.

Thus for our purpose we can propose a correspondence between the Q_{em} charge conservation and CPT-invariance, which, in our opinion, could have a geometrical origin in Poincaré duality [24]. This correspondence links in $D = 1 + 3$ space-time the matter–antimatter symmetry and Q_{em} charge conservation. This proposal can be interpreted as an attempt to find a correlation between external and internal symmetries, overcoming by this the Coleman–Mandula no-go theorem. In our conjecture, following to Poincaré duality, the internal properties of the particle fields could have a correlation/link with their space-time symmetry properties [24]. Of course, should be valid the reverse sentence: one can study space-time through the study of the particles properties.

In the beginning of the 90's, thanks to [12] and later to [13], the interest of the physics community was excited by the suggestion of uncompactified new extra dimensions and their link to high energy physics near the Standard Model energy scale. The main attraction of this approach is the possibility to reduce the mass Planck scale from 10^{19}GeV down to $\sim 1 - 10 \text{ TeV}$, an energy scale that can be explored experimentally in the LHC. So, a scenario with large or infinite uncompactified extra dimensions can be more acceptable than one based on a new physics near $\sim 10^{19} \text{ GeV}$.

Actually, to study the geometry of physical space-time with extra dimensions, one can proceed by investigating its symmetries. To build a quantum field theory in a space-time with extra dimensions ($D > 4$), some of which have a status different from our usual 4 ones, it seems inappropriate to use the usual the Lie–Killing–Cartan symmetries, as compactification, mentioned above, has taught us. So there could be just a chance to uncover new symmetries beyond the Lie–Cartan type. We propose to reach aim first by generalizing binary Lie algebras/symmetries to n -ary algebras/symmetries (*that is, instead of binary operations like addition or product of elements, one starts with composition laws involving at least n elements of the considered algebra*).

A mechanism of geometrical origin of electromagnetic charge Q_{em} and Dirac complex matter, baryon and lepton matter and baryon–antibaryon (lepton–antilepton) asymmetry, proton–electron non-stability etc, can be called Majorana–Diraco genesis and could be considered as a further step in the development of baryo–genesis and lepto–genesis . Note that baryo–genesis -theory or lepto–genesis -theory demands to explain the baryon or lepton asymmetry of the Universe some exotic particles, lepto–quarks or right–handed neutrino, correspondingly. The masses of such particles are very large: $M_{lq} \sim 10^{18}\text{GeV}$ and $M_{RH} \sim 10^{12} - 10^{14}\text{GeV}$, respectively. The Majorana–Diraco genesis is based on the existence of a new extra dimensional geometry and can predict the proton/electron decays at the TeV mass scale, which can be checked by modern experiments (LHC). Here one can find a similarity with the situation with $M_{Planck} \sim 10^{19}\text{GeV}$ in $D = 4$ and a possibility to diminish this parameter using some extra dimensions [12], [13].

The idea of Majorana–Diraco genesis starts from Dirac/Majorana equations and is related to the further attempts to solve the baryon asymmetry of the Universe linking such question with the origin Q_{em} charge symmetry. We would like to explore the idea of complexity of Dirac fermions and propose that the Dirac fermions in $D = 4$ have been produced from real fermions of higher dimensions by a complexification or by a mechanism unknown to us. There is one very remarkable fact

$$|Q_p + Q_e| < 10^{-21}, \quad (4)$$

which can indicate that the proton (quarks) and the electron can have a unique origin after the Big Bang .

This proposal suggests the existence in high-dimensional space of a class of real fermions connected with our fermions through a new interaction, which is based on a new symmetry beyond Lie, such an interaction predicts a non-stability of the electrons or protons. We can illustrate scale of such an interaction, taking for example two extra dimensions. To make estimations we are in the situation in which was E.Fermi before

the discovery the Yang-Mills interactions. So we will follow the idea he used for the construction of the four-fermion weak theory with the coupling constant GF , of dimension $[G_F] = [M^{-2}]$. So we can start from the multi-fermion $D = 6$ Fermi Lagrangian and the corresponding Fermi constant G_{Fs} , which could have dimension:

$$[G_{Fs}] = [M^{-p}], \quad p = 3, 4, 5, 6, \quad (5)$$

where p depends on the n -arity of a new interaction.

In our opinion, this dimension of coupling constant corresponds to a ternary gauge interaction, whose propagator could have a form like $[P(q)_p + M_s^p]^{-1}$, where $P(q)_p$ is a polynomial of the transferred momentum q of degree $p = 3, 4, \dots$. The form of such a propagator corresponds to a relativistic equation of the “gauge boson” flying in $D = 6$. We can propose that such a “gauge boson” will membrane origin, but not of string. So, the gauge variant of the Fermi $D = 6$ interactions could have a membrane origin and we can call them membrane gauge bosons. This is a completely new peculiarity of high-dimensional interactions based on ternary gauge symmetries. So for the tree-level calculations of the quark or charged-lepton decays into neutral real bulk fermions ν_S

$$q \mapsto m \nu_S, \quad e^\pm \mapsto m \nu_S, \quad (6)$$

taking the propagator of degree four one can get the following estimate for the partial width:

$$\Gamma(e \rightarrow nS) \sim g_S^4 \cdot \frac{m_{e/q}^9}{M_S^8}, \quad (7)$$

where $m_{e/q}$ is the electron mass and M_S is the membrane gauge boson mass.

To obtain the lower boundary for M_S one can compare the partial width for electron decay with the lifetime of the muon getting in D_4 -Fermi interactions. For the estimations one can take the following limit of the electron life-time: $\tau_e \leq 10^{25} \text{ years} \approx \pi \cdot 10^7 \cdot 10^{25} \text{ sec}$. This upper boundary gives the following estimate for

$$M_S \geq \left(\frac{g_2}{g_S}\right)^{1/2} \cdot 10 \cdot M_W. \quad (8)$$

From this, it is clear that the M_S can be also in the $\sim O(10) \text{ TeV}$ region. This mass scale is consistent with the upper boundary getting from the searching the proton decay. Note that in this approach one can study the proton decay problem in B - physics, in μ - and τ - decays also! There is also an interesting question about the spin structure of a new real bulk fermions, ν_S . Thank to the ternary symmetries they could have the $1/3$ spin structure. On algebraic language it means that a map of the internal ternary symmetry of new fermions into the symmetry of the ambient space-time is a nontrivial triple covering like in the case of the electron, its spin $1/2$ structure is related with the double covering maps: $SU(2) \mapsto SO(3)$ and/or $SL(2, C) \mapsto SO(3, 1)$ [7].

We started our discussion from the Majorana neutrino and step by step want to come to the idea of existence of completely new physics at the energy scale $\sim O(10)TeV$. Firstly, this physics could be related to the baryo-genesis problem and proton/electron non-stability, which we also have chances to observe in the $\mu-$, $\tau-$ lepton and B -meson rare decays. Secondly, since in our approach the Majorana neutrino “lives” in the higher-dimensional space-time describing by a new ternary symmetry generalized Lorentz symmetry we should also to change some fundamental principles of special theory of relativity, for example, the principle of maximal velocity. This one can check by measuring the neutrino velocity at high energies. If the energy scale $M_s \sim 10TeV$, we can expect an increase of the neutrino velocity $V/c - 1 \approx 10^{-4-5}$ at energies $E \sim 20 - 30GeV$ [26]. Thirdly, the Majorana-Dirac genesis gives us a chance to find the mechanism of creation of Dirac fermions and the $U(1)$ -electromagnetic vacuum of our Universe. This approach suggests the existence of a new fermion matter with some new interactions based on the ternary gauge symmetry. Thus this matter can be the candidate for the dark matter.

To understand much deeper the role of Majorana neutrinos in the study the ambient geometry of our Universe one can remind the role of photon and electron/antielectron in discovery of the Lorentz symmetry and Minkowsky $D = 3 + 1$ space-time. At the end of the XIX century the lasting efforts spent on the study of electromagnetism had been finished by Maxwell who wrote his fundamental equations [4] giving the theory of light. Lorentz found that the Maxwell equations are satisfied by the symmetry, which now bears his name. Poincaré extended the Lorentz symmetry by translations. Based on the Poincaré and Lorentz groups it was conjectured by Minkowski that our world is a 4-dimensional, Riemannian homogeneous space-time geometrical object with the structure described by the metrics $g_{\mu\nu} = diag(1, -1, -1, -1)$. Thus, one can say in short, that the study of electromagnetism-light extended our geometrical representations to the ambient structure of our world from dimension $D = 3$ to dimension $D = 4$.

Later, the theory of the electron just confirmed this conjecture. The spin structure of the electron can be described by a spinor wave function described by the $SU(2)$ group symmetry which is the double covering group of the $SO(3)$ group symmetry of the spatial part of our space. The extension of the external $SO(3)$ group to the Lorentz group $SO(1, 3)$, with its double covering group $SL(2, C)$, gave some additional degree of freedoms in the four-spinor dimensional spinor wave function which was used by Dirac to discover the antiparticles that were nicely confirmed by experiments. The theory of light and electron have been unified in quantum electrodynamics, based on the Poincaré /Lorentz symmetry and $U(1)_{em}$ gauge symmetry acting in the Minkowski $D=1+3$ space-time. One of the fundamental consequences of this quantum theory is the CPT theorem, which tells us that these two families of matter and antimatter should have some “degenerate” properties, *i.e.* the masses and life-times of particle and antiparticle must coincide. The ternary mass relation $m_{\nu_1} = m_{\nu_2} = m_{\nu_3}$ can be explained in analogy with two “generations” of particle and antiparticle in the binary Lorentz group, where the CPT theorem gives the exact relations between their masses: $m = \bar{m}$. If the origin of the three generations really is related to ternary symmetry, similar discrete symmetries could give such ternary mass relation for the masses of the three Majorana neutrinos living in $D = 6$. Since the Dirac fermions are living in $D = 4$ the ternary mass relation cannot be valid more. The

gauge invariance principle gave us the complete picture of the quantum theory of light which can be formulated in the language of the external and internal symmetry Poincaré duality:

$$SO(3,1) \leftrightarrow U(1)^{\text{em}} \quad (9)$$

Thus if we believe that we could understand the geometrical origin of $U(1)_{\text{em}}$ symmetry we must do the next step: to understand the geometrical origin of the SM , *i.e.* to understand the geometrical origin of the $SU(3^c)$, $SU(2_{\text{EW}}) \times U(1_Y)$ gauge symmetries and, consequently, to explain why the $SU(3^c)$ - symmetry is exact and the $SU(2_{\text{EW}}) \times U(1_Y)$ symmetry is broken?! The last comment is related with an alternative way to the Higgs mechanism of breaking the $SU(2) \times U(1)$ gauge symmetry to the $U(1)_{\text{em}}$ symmetry.

Our conjecture is that if the Majorana neutrinos living in the bulk can be described by a new ternary symmetry which is a generalization of the Lorentz symmetry. If the extra-dimensional space-time produce a new geometrical cycle the maximum speed limit for the high energetic neutrinos could be different from the velocity of the light and could be connected with the new fundamental constant of the second cycle, *i.e.* $C_2 \geq c_\nu \geq C_1 \equiv c$ [24, 15, 26]. Thus Majorana neutrinos can help us to go beyond the Lorentz symmetry and to see a new world through the new symmetries and new "gauge" interactions such as the membrane gauge bosons that can give an origin of a new "light" related with a new Abelian gauge symmetry, which would be called by "neutrino light" or "dark matter light" [25]:

1 Acknowledgements

I would like to thank particularly Tatiana Fabergé for her wonderful sense of the permanent help during the long period of working on this article.

The idea to write the article about the Majorana neutrinos was initiated by H.V. Klapdor-Kleingrothaus and I. Krivosheina, who invited me to the Gran-Sasso Laboratory on summer 2005, where I have got nice possibilities to study the intriguing results of the HEIDELBERG–MOSCOW experiment [1, 2, 3]. During seven last years we already discussed the possible role of Majorana neutrino in Majorana–Diraco genesis . The Majorana nature of neutrino is playing the crucial role in this scheme. This idea was closely related with the possible new geometrical properties ($D > 4$) of Majorana neutrino to check which we suggested in measuring the neutrino velocity at high energies [15]. The progress what made in the construction of the new neutrino detectors in Gran Sasso Underground Laboratory gives us the independent possibilities to check is neutrino of Majorana nature or not [15]. This question is dramatically difficult both in theory and in experiment. (The neutrino short base experiment to study if the neutrino is a tachyon or not, was done in FNAL in the 1977.) The other possibility to check the Majorana nature of neutrino can be related with the proton decay problem. In the scheme of Majorana–Diraco genesis the proton non-stability problem could be solved also in the $e-$, $\mu-$, $\tau-$ lepton and $b-$ meson rare decay experiments. The Majorana nature of the neutrino gave a lot of new and non-ordinary ideas for the probing of which I would like to express my

thanks to L. Alvarez-Gaumé, U. Aglietti, V. Ammosov, I. Antoniadis, P. Aurenche, M. Baldo-Ceolin, G. Bélanger, G. Costa, P. Chankowski, A. De Rujula, A. Dubrovskiy, J. Ellis, L. Fellin, H. Klapdor-Kleingrothaus, I. Krivosheina, L. Lipatov, A. Liparteliani, A. Masiero, R. Obergfeld, M. Ruggier, A. Sabio-Vera, J-B. Zuber. It is pleasure for me to thank Z.Berezhiani and the participants of the X Summer Institute 2005 and XI Summer Institute 2006 in Gran-Sasso for the nice possibilities to discuss the ideas of this work.

I also thank very much G. Girardi for reading manuscript and many comments.

References

- [1] H.V. Klapdor-Kleingrothaus et al., *Mod. Phys. Lett. A* **16** (2001) 2409–2420 and *Found. Phys.* **31** (2002) 1181–1123 (*Corrigenda*, 2003).
- [2] H.V. Klapdor-Kleingrothaus et al., *Phys. Lett. B* **586** (2004) 198–212.
- [3] H.V. Klapdor-Kleingrothaus et al., *Nucl. Instrum. Methods A* **522** (2004) 371–406.
- [4] J.C. Maxwell, *An elementary treatise on electricity*, 2nd ed. (Clarendon Press, Oxford, 1988).
- [5] P.A.M. Dirac, “The quantum theory of electron”, *Proc. Roy. Soc.* **A117** (1928) 610.
- [6] E. Majorana, “Teoria simmetrica dell’ellectrone e del positrone”, *Nuovo Cimento* **14** (1937) 171.
- [7] C. Chevalley, *The algebraic theory of spinors* (Columbia University Press, New York, 1954).
- [8] M. Raush de Traubenberg, “Clifford algebra in physics”, hep-th/0506011 (2005) and “Algèbres de Clifford supersymétrie et symétrie Z_n – Applications en théorie des champs”, Orsay report LPT 97-02 (1997).
- [9] A. Dubrovsky. and G. Volkov, *Ternary numbers and algebras. Reflexive numbers and Berger graphs* To be published (2006).
- [10] N. Bogoljubov and D. Shirkov, *Introduction to quantum field theory*.
- [11] A. Capelli, C. Itzykson and J.-B. Zuber, *Commun. Math.phys* **184** (1987), 1-26, MR **89b**, 81178.
- [12] I. Antoniadis, *Phys. Lett. B* **246** (1990) 377.
- [13] N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett. B* **429** (1998) 263;
I. Antoniadis, N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys. Lett. B* **436** (1998) 256.
- [14] V. Rubakov, “Large and infinite extra dimensions”, *Phys. Usp.* **44** (2001) 871.

- [15] V. Ammosov and G. Volkov, “Can neutrino probe extra dimension”, Padua preprint DFPD-00/TH/39, hep-ph/008 (2000).
- [16] Y. Nambu, “Generalized Hamiltonian mechanics”, *Phys. Rev.* **D7** (1973) 2405–2412.
- [17] V.T. Filippov, “ n -ary Lie algebras”, *Sibirskii Matematicheskii Zhurnal* **26**, 6 (1985) 126–140.
- [18] R. Kerner, “Ternary algebraic structures and their applications in physics”, math-ph/0011023, *Proc. Conference ICGTMP “Group-23”*, Dubna, 2000.
- [19] M. Dubois-Violette, “Generalized differential spaces with $d^N = 0$ and the q -differential calculus”, q-alg/9609012 (1996).
- [20] V. Abramov, R. Kerner, B. Le Roy, *Hypersymmetry: a Z_3 -graded generalization of supersymmetry* J.Math.Phys. 38 (1997) 1650-1669, hep-th/9607143
V. Abramov, *Z_3 -graded analogues of Clifford algebras and generalization of supersymmetry* Lecture given at XXI Colloquium on Group Theoretical Methods in Physics, Goslar, 1996, hep-th/9607241
- [21] F. Anselmo, J. Ellis, D. Nanopoulos and G. Volkov, “Towards an algebraic classification of Calabi–Yau Manifolds: Study of K3 spaces”, *Phys. Part. Nucl.* **32** (2001) 318–375, and *Fiz. Elem. Chast. Atom. Yadra* **32** (2001) 605–698.
- [22] G. Volkov, “Hunting for the new symmetries in Calabi–Yau jungles”, *Int. J. Mod. Phys.* **A19** (2004) 4835–4860, hep-th/0402042.
- [23] L. Lipatov, A. Sabio-Vera, V. Velighanin and G. Volkov, *Reflexive numbers and Berger graphs from Calabi-Yau spaces* hep-th/0501101 (2005) to be published in IJMP.
- [24] G. Volkov, “From Galois theory to the ultra-radical high energy physics”, preprint LAPTH-958-02 (2002).
- [25] www.neutrinolight.org
- [26] G. Volkov, talk at the OPERA Meeting, 10 January 2006.